

Logic in Action

Chapter 2: Propositional Logic

`http://www.logicinaction.org/`

Example (1)

In a restaurant, your Father has ordered Fish, your Mother ordered Vegetarian, and you ordered Meat. Out of the kitchen comes some new person carrying the three plates. What will happen?

Example (2)

Three guests are sitting at a table. The waitress asks: **“Does everyone want coffee”**. The first guest says: **“I don’t know”**. The second guest now says: **“I don’t know”**. Then the third guest says: **“No, not everyone wants coffee”**. The waitress comes back and gives the right people their coffees. Assuming that at the beginning each guest only knows about himself, which was the waitress reasoning? Who gets coffee and who does not?

Example (3)

1	.	.
.	.	2
.	.	.

Example (3)

1	·	3
·	·	2
·	·	·

Example (3)

1	2	3
.	.	2
.	.	1

Example (3)

1	2	3
3	.	2
.	3	1

Example (3)

1	2	3
3	1	2
2	3	1

Example (4)

If you take the medication, you will get better.

You are taking the medication.

?

So, you will get better.

Example (4)

If you take the medication, you will get better.

✓ You are taking the medication.

So, you will get better.

Example (4)

If you take the medication, you will get better.

You are taking the medication.



So, you will get better.

If you take the medication, you will get better.

You are getting better.

?

So, you took the medication.

Example (4)

If you take the medication, you will get better.

You are taking the medication.



So, you will get better.

If you take the medication, you will get better.

You are getting better.



So, you took the medication.

Example (5)

If you take the medication, you will get better.

But you are not taking the medication.

?

So, you will not get better.

Example (5)

If you take the medication, you will get better.

But you are not taking the medication.

X

So, you will not get better.

Example (5)

x

If you take the medication, you will get better.
But you are not taking the medication.

So, you will not get better.

?

If you take the medication, you will get better.
But you are not getting better.

So, you have not taken the medication.

Example (5)

If you take the medication, you will get better.

But you are not taking the medication.

✗

So, you will not get better.

If you take the medication, you will get better.

But you are not getting better.

✓

So, you have not taken the medication.

Valid inference

$$\frac{A_1, \dots, A_n}{C}$$

Valid inference

$$\frac{A_1, \dots, A_n}{C}$$

An inference is **valid** if and only if **every** time **all** the premises are true, the conclusion is also true.

What a valid inference tells us?

Suppose the following inference is valid

$$\frac{A_1, \dots, A_n}{C}$$

Then

What a valid inference tells us?

Suppose the following inference is valid

$$\frac{A_1, \dots, A_n}{C}$$

Then

- 1 if **all** the premises A_1, \dots, A_n are true, so is the conclusion C .

What a valid inference tells us?

Suppose the following inference is valid

$$\frac{A_1, \dots, A_n}{C}$$

Then

- 1 if **all** the premises A_1, \dots, A_n are true, so is the conclusion C .
- 2 if the conclusion C is false, **at least one** premise A_i is false.

Looking for patterns (1)

Two valid inferences:

If you take the medication, then you will get better.

You are taking the medication.

So, you will get better.

If you jump from a 4th floor, then you will fly.

You jump from a 4th floor.

So, you will fly.

Looking for patterns (1)

Two valid inferences:

If **you take the medication**, then **you will get better**.
You are taking the medication.

So, **you will get better**.

If **you jump from a 4th floor**, then **you will fly**.
You jump from a 4th floor.

So, **you will fly**.

Looking for patterns (1)

Two valid inferences:

If A , then B .

A .

So, B .

If E , then F .

E .

So, F .

Looking for patterns (2)

Another valid inference:

If you take the medication, then you will get better.

You are not getting better.

So, you are not taking the medication.

Looking for patterns (2)

Another valid inference:

If **you take the medication**, then **you will get better**.

You are not getting better.

So, **you are not taking the medication.**

Looking for patterns (2)

Another valid inference:

If A , then B .

not B .

So, not A .

Looking for patterns (3)

And yet another:

An integer x is even or odd.

If x is even, then $x + x$ is even.

If x is odd, then $x + x$ is even.

So, $x + x$ is even.

Looking for patterns (3)

And yet another:

An integer x is **even** or **odd**.

If x is **even**, then $x + x$ is **even**.

If x is **odd**, then $x + x$ is **even**.

So, $x + x$ is **even**.

Looking for patterns (3)

And yet another:

An integer x is A_1 or A_2 .

If x is A_1 , then C .

If x is A_2 , then C .

So, C .

Looking for patterns (3)

And yet another:

An integer x is A_1 or A_2 .

If x is A_1 , then C .

If x is A_2 , then C .

So, C .

Can you think of others?

The main question

How can we recognize valid inference patterns?

Example (6)



- A1 At least one of them is guilty.
- A2 Not all of them are guilty.
- A3 If Mrs White is guilty, then Colonel Mustard helped her (he is guilty too).
- A4 If Miss Scarlet is innocent then so is Colonel Mustard.

Example (6)



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

A1 At least one of them is guilty.



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

A1 At least one of them is guilty.



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

A1 At least one of them is guilty.

A2 Not all of them are guilty.



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

A1 At least one of them is guilty.

A2 Not all of them are guilty.



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

- A1 At least one of them is guilty.
- A2 Not all of them are guilty.
- A3 If Mrs White is guilty, then Colonel Mustard helped her (he is guilty too).



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

- A1 At least one of them is guilty.
- A2 Not all of them are guilty.
- A3 If Mrs White is guilty, then Colonel Mustard helped her (he is guilty too).



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

- A1 At least one of them is guilty.
- A2 Not all of them are guilty.
- A3 If Mrs White is guilty, then Colonel Mustard helped her (he is guilty too).
- A4 If Miss Scarlet is innocent then so is Colonel Mustard.



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

- A1 At least one of them is guilty.
- A2 Not all of them are guilty.
- A3 If Mrs White is guilty, then Colonel Mustard helped her (he is guilty too).
- A4 If Miss Scarlet is innocent then so is Colonel Mustard.



innocent	innocent	innocent
innocent	innocent	guilty
innocent	guilty	innocent
innocent	guilty	guilty
guilty	innocent	innocent
guilty	innocent	guilty
guilty	guilty	innocent
guilty	guilty	guilty

Example (6)

$$? \frac{A_1, A_2, A_3, A_4}{\text{Miss Scarlet is guilty}}$$

Example (6)

✓ $\frac{A_1, A_2, A_3, A_4}{\text{Miss Scarlet is guilty}}$

Example (6)

$$\checkmark \frac{A_1, A_2, A_3, A_4}{\text{Miss Scarlet is guilty}}$$

In **every** situation in which A_1, A_2, A_3 and A_4 are all true, “**Miss Scarlet is guilty**” is true.

Example (6)

$$\checkmark \frac{A_1, A_2, A_3, A_4}{\text{Miss Scarlet is guilty}}$$

In **every** situation in which A_1, A_2, A_3 and A_4 are all true, “**Miss Scarlet is guilty**” is true.

$$? \frac{A_1, A_2, A_3, A_4}{\text{Mrs White is innocent}}$$

Example (6)

$$\checkmark \frac{A_1, A_2, A_3, A_4}{\text{Miss Scarlet is guilty}}$$

In **every** situation in which A_1, A_2, A_3 and A_4 are all true, “**Miss Scarlet is guilty**” is true.

$$\checkmark \frac{A_1, A_2, A_3, A_4}{\text{Mrs White is innocent}}$$

Example (6)

$$\checkmark \frac{A_1, A_2, A_3, A_4}{\text{Miss Scarlet is guilty}}$$

In **every** situation in which A_1, A_2, A_3 and A_4 are all true, “**Miss Scarlet is guilty**” is true.

$$\checkmark \frac{A_1, A_2, A_3, A_4}{\text{Mrs White is innocent}}$$

In **every** situation in which A_1, A_2, A_3 and A_4 are all true, “**Mrs White is innocent**” is true.

Example (6)

$$? \frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is guilty}}$$

Example (6)

$$\times \frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is guilty}}$$

Example (6)

$$\times \frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is guilty}}$$

There is a situation in which A_1 , A_2 , A_3 and A_4 are all true, but “Colonel Mustard is guilty” is false (there is a counter-example).

Example (6)

\times $\frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is guilty}}$

There is a situation in which A_1, A_2, A_3 and A_4 are all true, but “Colonel Mustard is guilty” is false (there is a counter-example).

? $\frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is innocent}}$

Example (6)

$$\times \frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is guilty}}$$

There is a situation in which A_1, A_2, A_3 and A_4 are all true, but “Colonel Mustard is guilty” is false (there is a **counter-example**).

$$\times \frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is innocent}}$$

Example (6)

$$\times \frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is guilty}}$$

There is a situation in which A_1, A_2, A_3 and A_4 are all true, but “Colonel Mustard is guilty” is false (there is a counter-example).

$$\times \frac{A_1, A_2, A_3, A_4}{\text{Colonel Mustard is innocent}}$$

There is a situation in which A_1, A_2, A_3 and A_4 are all true, but “Colonel Mustard is innocent” is false (there is a counter-example).

Ingredients of the **propositional** language

Ingredients of the **propositional** language

- ① Basic (*atomic*) statements (**propositions**):

$$p, q, r, \dots$$

Ingredients of the **propositional** language

- ① Basic (*atomic*) statements (**propositions**):

$$p, q, r, \dots$$

- ② Operators to build more statements:

“ not ...”	becomes	$\neg \dots$
“... and ...”	becomes	$\dots \wedge \dots$
“... or ...”	becomes	$\dots \vee \dots$
“ if ... then ”	becomes	$\dots \rightarrow \dots$
“... if and only if ...”	becomes	$\dots \leftrightarrow \dots$

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

$$\neg\varphi \in \mathcal{L}_P,$$

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

$$\neg\varphi \in \mathcal{L}_P, \quad (\varphi \wedge \psi) \in \mathcal{L}_P,$$

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

$$\begin{aligned} \neg\varphi &\in \mathcal{L}_P, & (\varphi \wedge \psi) &\in \mathcal{L}_P, \\ (\varphi \vee \psi) &\in \mathcal{L}_P, \end{aligned}$$

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

$$\begin{aligned} \neg\varphi \in \mathcal{L}_P, & \quad (\varphi \wedge \psi) \in \mathcal{L}_P, & \quad (\varphi \rightarrow \psi) \in \mathcal{L}_P, \\ & \quad (\varphi \vee \psi) \in \mathcal{L}_P, \end{aligned}$$

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

$$\begin{array}{lll} \neg\varphi \in \mathcal{L}_P, & (\varphi \wedge \psi) \in \mathcal{L}_P, & (\varphi \rightarrow \psi) \in \mathcal{L}_P, \\ (\varphi \vee \psi) \in \mathcal{L}_P, & & (\varphi \leftrightarrow \psi) \in \mathcal{L}_P. \end{array}$$

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

$$\begin{array}{lll} \neg\varphi \in \mathcal{L}_P, & (\varphi \wedge \psi) \in \mathcal{L}_P, & (\varphi \rightarrow \psi) \in \mathcal{L}_P, \\ (\varphi \vee \psi) \in \mathcal{L}_P, & & (\varphi \leftrightarrow \psi) \in \mathcal{L}_P. \end{array}$$

- 3 Nothing else is in \mathcal{L}_P .

The propositional language

The **language** \mathcal{L}_P is a set of formulas satisfying:

- 1 All the basic propositions are in \mathcal{L}_P :

$$p \in \mathcal{L}_P, \quad q \in \mathcal{L}_P, \quad r \in \mathcal{L}_P, \quad \dots$$

- 2 If $\varphi \in \mathcal{L}_P$ and $\psi \in \mathcal{L}_P$, then

$$\begin{array}{lll} \neg\varphi \in \mathcal{L}_P, & (\varphi \wedge \psi) \in \mathcal{L}_P, & (\varphi \rightarrow \psi) \in \mathcal{L}_P, \\ (\varphi \vee \psi) \in \mathcal{L}_P, & & (\varphi \leftrightarrow \psi) \in \mathcal{L}_P. \end{array}$$

- 3 Nothing else is in \mathcal{L}_P .

In practice, we will avoid parenthesis if they are not necessary.

Constructing formulas

The construction of a formula can be seen as building a **tree**.

Constructing formulas

The construction of a formula can be seen as building a **tree**.

$$((\neg p \vee q) \rightarrow r)$$


Constructing formulas

The construction of a formula can be seen as building a **tree**.

$$\begin{array}{c} ((\neg p \vee q) \rightarrow r) \\ | \\ \rightarrow \end{array}$$


Constructing formulas

The construction of a formula can be seen as building a **tree**.




Constructing formulas

The construction of a formula can be seen as building a **tree**.




Constructing formulas

The construction of a formula can be seen as building a **tree**.




Constructing formulas

The construction of a formula can be seen as building a **tree**.




Constructing formulas

The construction of a formula can be seen as building a **tree**.




Constructing formulas

The construction of a formula can be seen as building a **tree**.




Constructing formulas

The construction of a formula can be seen as building a **tree**.





Constructing formulas

The construction of a formula can be seen as building a **tree**.





Constructing formulas

The construction of a formula can be seen as building a **tree**.





Constructing formulas

The construction of a formula can be seen as building a **tree**.





Constructing formulas

The construction of a formula can be seen as building a **tree**.



Constructing formulas

The construction of a formula can be seen as building a **tree**.



Evaluating formulas

How do we know if a given formula φ is **true** or **false**?

Evaluating formulas

How do we know if a given formula φ is **true** or **false**?

- We need the **truth-values** of the basic propositions p, q, r, \dots that appear in φ .

Evaluating formulas

How do we know if a given formula φ is **true** or **false**?

- We need the **truth-values** of the basic propositions p, q, r, \dots that appear in φ .
- We need to know the **meaning** of $\neg, \wedge, \vee, \rightarrow$ and \leftrightarrow .

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
-----------	---------------

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
1	0
0	1

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
1	0
0	1

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
1	0
0	1

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
1	0
0	1

or, in a shorter format:

\neg	φ

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
1	0
0	1

or, in a shorter format:

\neg	φ
	1
	0

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
1	0
0	1

or, in a shorter format:

\neg	φ
0	1
	0

Behaviour of the connectives (1)

Use 1 for **true**, and 0 for **false**.

For **negation** \neg

φ	$\neg\varphi$
1	0
0	1

or, in a shorter format:

\neg	φ
0	1
1	0

Behaviour of the connectives (2)

For **conjunction** \wedge

$$\frac{\varphi \quad \wedge \quad \psi}{}$$

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1		1
1		0
0		1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1		0
0		1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0		1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0	0	0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0	0	0

For **disjunction** \vee

φ	\vee	ψ
1	1	1
1	1	0
0	1	1
0	0	0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0	0	0

For **disjunction** \vee

φ	\vee	ψ
1		1
1		0
0		1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0	0	0

For **disjunction** \vee

φ	\vee	ψ
1	1	1
1		0
0		1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0	0	0

For **disjunction** \vee

φ	\vee	ψ
1	1	1
1	1	0
0		1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0	0	0

For **disjunction** \vee

φ	\vee	ψ
1	1	1
1	1	0
0	1	1
0		0

Behaviour of the connectives (2)

For **conjunction** \wedge

φ	\wedge	ψ
1	1	1
1	0	0
0	0	1
0	0	0

For **disjunction** \vee

φ	\vee	ψ
1	1	1
1	1	0
0	1	1
0	0	0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

$$\frac{\varphi \leftrightarrow \psi}{\quad}$$

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1		1
1		0
0		1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1		0
0		1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0		1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0	1	0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0	1	0

For **implication** \rightarrow

φ	\rightarrow	ψ
1		1
1		0
0		1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0	1	0

For **implication** \rightarrow

φ	\rightarrow	ψ
1		1
1		0
0		1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0	1	0

For **implication** \rightarrow

φ	\rightarrow	ψ
1	1	1
1		0
0		1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0	1	0

For **implication** \rightarrow

φ	\rightarrow	ψ
1	1	1
1	0	0
0		1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0	1	0

For **implication** \rightarrow

φ	\rightarrow	ψ
1	1	1
1	0	0
0	1	1
0		0

Behaviour of the connectives (3)

For **equivalence** \leftrightarrow

φ	\leftrightarrow	ψ
1	1	1
1	0	0
0	0	1
0	1	0

For **implication** \rightarrow

φ	\rightarrow	ψ
1	1	1
1	0	0
0	1	1
0	1	0

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Example: assume $P = \{p, q\}$.

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Example: assume $P = \{p, q\}$.

There are **four** different valuations (**four** different situations):

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Example: assume $P = \{p, q\}$.

There are **four** different valuations (**four** different situations):

$$\overline{\overline{V_1(p) = 1 \quad V_1(q) = 1}}$$

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Example: assume $P = \{p, q\}$.

There are **four** different valuations (**four** different situations):

$V_1(p) = 1$	$V_1(q) = 1$
$V_2(p) = 1$	$V_2(q) = 0$

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Example: assume $P = \{p, q\}$.

There are **four** different valuations (**four** different situations):

$V_1(p) = 1$	$V_1(q) = 1$
$V_2(p) = 1$	$V_2(q) = 0$
$V_3(p) = 0$	$V_3(q) = 1$

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Example: assume $P = \{p, q\}$.

There are **four** different valuations (**four** different situations):

$V_1(p) = 1$	$V_1(q) = 1$
$V_2(p) = 1$	$V_2(q) = 0$
$V_3(p) = 0$	$V_3(q) = 1$
$V_4(p) = 0$	$V_4(q) = 0$

Valuations

Valuation. Let $P = \{p, q, r, \dots\}$ be a set of atomic propositions. A **valuation** V from P to $\{0, 1\}$ assigns to each element of P a unique truth-value.

Example: assume $P = \{p, q\}$.

There are **four** different valuations (**four** different situations):

$V_1(p) = 1$	$V_1(q) = 1$
$V_2(p) = 1$	$V_2(q) = 0$
$V_3(p) = 0$	$V_3(q) = 1$
$V_4(p) = 0$	$V_4(q) = 0$

How many for $P = \{p\}$? How many for $P = \{p, q, r\}$?

Evaluating formulas in one situation

$$(\neg p) \wedge q \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \quad (\neg \quad p) \quad \wedge \quad q \quad \boxed{}$$

Evaluating formulas in one situation

$V:$	$(\neg$	$p)$	\wedge	q	<input type="text"/>
	1	0		1	

Evaluating formulas in one situation

$V:$	$(\neg$	$p)$	\wedge	q	$\boxed{}$
	1	0	1	1	

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & \mathbf{1} & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

Evaluating formulas in one situation

$$V: \quad \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$(p \wedge (p \rightarrow q)) \rightarrow q \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & & 1 & \rightarrow & 0 & \rightarrow & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & \mathbf{1} & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & & 1 & 0 & 0 & & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \quad \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \quad \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & \rightarrow & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{ccccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & \mathbf{1} & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & \mathbf{1} & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$\neg \quad \neg \quad p \quad \boxed{\phantom{\text{truth value}}}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ & & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ & 1 & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ 0 & 1 & 0 \end{array} \quad \boxed{\phantom{V \models \text{formula}}}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{ccccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ 0 & 1 & 0 \end{array} \quad \boxed{V \not\models \neg\neg p}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ 0 & 1 & 0 \end{array} \quad \boxed{V \not\models \neg\neg p}$$

$$(p \rightarrow q) \vee (q \rightarrow p) \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ 0 & 1 & 0 \end{array} \quad \boxed{V \not\models \neg\neg p}$$

$$V: \begin{array}{cccc} (p & \rightarrow & q) & \vee & (q & \rightarrow & p) \\ 0 & & 1 & & 1 & & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ 0 & 1 & 0 \end{array} \quad \boxed{V \not\models \neg\neg p}$$

$$V: \begin{array}{ccccc} (p & \rightarrow & q) & \vee & (q & \rightarrow & p) \\ 0 & 1 & 1 & & 1 & & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ 0 & 1 & 0 \end{array} \quad \boxed{V \not\models \neg\neg p}$$

$$V: \begin{array}{cccc} (p & \rightarrow & q) & \vee & (q & \rightarrow & p) \\ 0 & 1 & 1 & & 1 & 0 & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & 1 & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & 1 & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ 0 & 1 & 0 \end{array} \quad \boxed{V \not\models \neg\neg p}$$

$$V: \begin{array}{cccc} (p & \rightarrow & q) & \vee & (q & \rightarrow & p) \\ 0 & 1 & 1 & 1 & 1 & 0 & 0 \end{array} \quad \boxed{}$$

Evaluating formulas in one situation

$$V: \begin{array}{cccc} (\neg & p) & \wedge & q \\ 1 & 0 & \mathbf{1} & 1 \end{array} \quad \boxed{V \models (\neg p) \wedge q}$$

$$V: \begin{array}{cccccc} (p & \wedge & (p & \rightarrow & q)) & \rightarrow & q \\ 1 & 0 & 1 & 0 & 0 & \mathbf{1} & 0 \end{array} \quad \boxed{V \models (p \wedge (p \rightarrow q)) \rightarrow q}$$

$$V: \begin{array}{ccc} \neg & \neg & p \\ \mathbf{0} & 1 & 0 \end{array} \quad \boxed{V \not\models \neg\neg p}$$

$$V: \begin{array}{cccc} (p & \rightarrow & q) & \vee & (q & \rightarrow & p) \\ 0 & 1 & 1 & \mathbf{1} & 1 & 0 & 0 \end{array} \quad \boxed{V \models (p \rightarrow q) \vee (q \rightarrow p)}$$

Evaluating formulas in all possible situations

$$(p \wedge (p \rightarrow q)) \rightarrow q$$

Evaluating formulas in all possible situations

p	\wedge	$(p \rightarrow q)$	\rightarrow	q
1		1		1
1		1		0
0		0		1
0		0		0

Evaluating formulas in all possible situations

p	\wedge	$(p \rightarrow q)$	\rightarrow	q
1	1	1	1	1
1	1	0	0	0
0	0	1	1	1
0	0	1	0	0

Evaluating formulas in all possible situations

p	\wedge	$(p \rightarrow q)$	\rightarrow	q
1	1	1	1	1
1	0	1	0	0
0	0	0	1	1
0	0	0	1	0

Evaluating formulas in all possible situations

p	\wedge	$(p \rightarrow q)$	\rightarrow	q		
1	1	1	1	1	1	1
1	0	1	0	0	1	0
0	0	0	1	1	1	1
0	0	0	1	0	1	0

Evaluating formulas in all possible situations

p	\wedge	$(p \rightarrow q)$	\rightarrow	q		
1	1	1	1	1	1	1
1	0	1	0	0	1	0
0	0	0	1	1	1	1
0	0	0	1	0	1	0

 $\neg \neg p$

Evaluating formulas in all possible situations

p	\wedge	$(p \rightarrow q)$	\rightarrow	q		
1	1	1	1	1	1	1
1	0	1	0	0	1	0
0	0	0	1	1	1	1
0	0	0	1	0	1	0

\neg	\neg	p
		1
		0

Evaluating formulas in all possible situations

p	\wedge	$(p \rightarrow q)$	\rightarrow	q
1	1	1	1	1
1	0	1	0	0
0	0	0	1	1
0	0	0	1	0

\neg	\neg	p
0	1	
1	0	

Evaluating formulas in all possible situations

$(p \wedge (p \rightarrow q)) \rightarrow q$	q
1	1
1	0
0	1
0	0

\neg	\neg	p
1	0	1
0	1	0

Classification of formulas according to their behaviour

Classification of formulas according to their behaviour

- Those that are never true (**contradiction**):

$$p \wedge (\neg p), \dots$$

Classification of formulas according to their behaviour

- Those that are never true (**contradiction**):

$$p \wedge (\neg p), \dots$$

- Those that can be true (**satisfiable**):

$$(\neg p) \vee q, \dots$$

Classification of formulas according to their behaviour

- Those that are never true (**contradiction**):

$$p \wedge (\neg p), \dots$$

- Those that can be true (**satisfiable**):

$$(\neg p) \vee q, \dots$$

- Those that are always true (**valid**, **tautology**):

$$(p \wedge (p \rightarrow q)) \rightarrow q, \dots$$

If the formula φ is valid, we write $\models \varphi$

Valid inference

$$\text{Inference: } \frac{\varphi_1, \dots, \varphi_n}{\psi}$$

Valid inference

$$\text{Inference: } \frac{\varphi_1, \dots, \varphi_n}{\psi}$$

Valid inference. An inference is **valid** if and only if **every** time (**every** situation) in which **all** premises $\varphi_1, \dots, \varphi_n$ are true, ψ is also true.

Valid inference

$$\text{Inference: } \frac{\varphi_1, \dots, \varphi_n}{\psi}$$

Valid inference. An inference is **valid** if and only if **every** time (**every** situation) in which **all** premises $\varphi_1, \dots, \varphi_n$ are true, ψ is also true.

We also say ψ is a **logical consequence** of $\varphi_1, \dots, \varphi_n$.

Valid inference

$$\text{Inference: } \frac{\varphi_1, \dots, \varphi_n}{\psi}$$

Valid inference. An inference is **valid** if and only if **every** time (**every** situation) in which **all** premises $\varphi_1, \dots, \varphi_n$ are true, ψ is also true.

We also say ψ is a **logical consequence** of $\varphi_1, \dots, \varphi_n$.

We will write $\varphi_1, \dots, \varphi_n \models \psi$

Examples

Our previous patterns:

$$\overline{\overline{p \mid (p \rightarrow q) \parallel q}}$$

Examples

Our previous patterns:

p	$(p \rightarrow q)$	q
1	1	1
1	0	0
0	1	1
0	1	0

Examples

Our previous patterns:

	p	$(p \rightarrow q)$	q
\rightarrow	1	1	1
	1	0	0
	0	1	1
	0	1	0

Examples

Our previous patterns:

	p	$(p \rightarrow q)$	q
\rightarrow	1	1	1
	1	0	0
	0	1	1
	0	1	0

$\neg q$	$(p \rightarrow q)$	$\neg p$

Examples

Our previous patterns:

	p	$(p \rightarrow q)$	q
\rightarrow	1	1	1
	1	0	0
	0	1	1
	0	1	0

	$\neg q$	$(p \rightarrow q)$	$\neg p$
	0	1	0
	1	0	0
	0	1	1
	1	0	1

Examples

Our previous patterns:

	p	$(p \rightarrow q)$	q
\rightarrow	1	1	1
	1	0	0
	0	1	1
	0	1	0

	$\neg q$	$(p \rightarrow q)$	$\neg p$
	0	1	0
	1	0	0
	0	1	1
\rightarrow	1	0	1

Examples

Our previous patterns:

	p	$(p \rightarrow q)$	q
\rightarrow	1	1	1
	1	0	0
	0	1	1
	0	1	0

	$\neg q$	$(p \rightarrow q)$	$\neg p$
	0	1	0
	1	0	0
	0	1	1
\rightarrow	1	0	1

What about the others?

Further definitions

Further definitions

- Two formulas φ and ψ are **logically equivalent** ($\varphi \equiv \psi$) if and only if $\varphi \models \psi$ and $\psi \models \varphi$.

Further definitions

- Two formulas φ and ψ are **logically equivalent** ($\varphi \equiv \psi$) if and only if $\varphi \models \psi$ and $\psi \models \varphi$.
- A set of formulas X is **satisfiable** if and only if **there is one valuation** that makes **every** formula in X true.

Symbolic inference

Symbolic inference

- A **proof** is a finite sequence of formulas where each formula is either an *axiom* or else it has been inferred from previous formulas by using an **inference rule**.

Symbolic inference

- A **proof** is a finite sequence of formulas where each formula is either an *axiom* or else it has been inferred from previous formulas by using an **inference rule**.
- A formula is a **theorem** if it occurs in a proof.

Symbolic inference

- A **proof** is a finite sequence of formulas where each formula is either an *axiom* or else it has been inferred from previous formulas by using an **inference rule**.
- A formula is a **theorem** if it occurs in a proof.
- A set of axioms and rules is called an **axiom system** or an **axiomatization**.

Symbolic inference

- A **proof** is a finite sequence of formulas where each formula is either an *axiom* or else it has been inferred from previous formulas by using an **inference rule**.
- A formula is a **theorem** if it occurs in a proof.
- A set of axioms and rules is called an **axiom system** or an **axiomatization**.
 - An axiom system is **sound** for a logic if every theorem is valid in the logic.

Symbolic inference

- A **proof** is a finite sequence of formulas where each formula is either an *axiom* or else it has been inferred from previous formulas by using an **inference rule**.
- A formula is a **theorem** if it occurs in a proof.
- A set of axioms and rules is called an **axiom system** or an **axiomatization**.
 - An axiom system is **sound** for a logic if every theorem is valid in the logic.
 - An axiom system is **complete** if every valid formula of the logic is a theorems.

Proof system

The following axiom system is sound and complete for propositional logic:

Proof system

The following axiom system is sound and complete for propositional logic:

① $(\varphi \rightarrow (\psi \rightarrow \varphi))$.

Proof system

The following axiom system is sound and complete for propositional logic:

- 1 $(\varphi \rightarrow (\psi \rightarrow \varphi))$.
- 2 $((\varphi \rightarrow (\psi \rightarrow \chi)) \rightarrow ((\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \chi)))$.

Proof system

The following axiom system is sound and complete for propositional logic:

- 1 $(\varphi \rightarrow (\psi \rightarrow \varphi))$.
- 2 $((\varphi \rightarrow (\psi \rightarrow \chi)) \rightarrow ((\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \chi)))$.
- 3 $((\neg\varphi \rightarrow \neg\psi) \rightarrow (\psi \rightarrow \varphi))$.

Proof system

The following axiom system is sound and complete for propositional logic:

- 1 $(\varphi \rightarrow (\psi \rightarrow \varphi))$.
- 2 $((\varphi \rightarrow (\psi \rightarrow \chi)) \rightarrow ((\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \chi)))$.
- 3 $((\neg\varphi \rightarrow \neg\psi) \rightarrow (\psi \rightarrow \varphi))$.
- 4 **Modus ponens (MP)**: from φ and $\varphi \rightarrow \psi$, infer ψ .

Example

Example

1. $p \rightarrow ((q \rightarrow p) \rightarrow p)$ Instance of axiom 1

Example

1. $p \rightarrow ((q \rightarrow p) \rightarrow p)$ Instance of axiom 1
2. $(p \rightarrow ((q \rightarrow p) \rightarrow p)) \rightarrow ((p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p))$ Instance of axiom 2

Example

1. $p \rightarrow ((q \rightarrow p) \rightarrow p)$ Instance of axiom 1
2. $(p \rightarrow ((q \rightarrow p) \rightarrow p)) \rightarrow ((p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p))$ Instance of axiom 2
3. $(p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p)$ MP from steps 1 and 2

Example

- | | | |
|----|---|-----------------------|
| 1. | $p \rightarrow ((q \rightarrow p) \rightarrow p)$ | Instance of axiom 1 |
| 2. | $(p \rightarrow ((q \rightarrow p) \rightarrow p)) \rightarrow ((p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p))$ | Instance of axiom 2 |
| 3. | $(p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p)$ | MP from steps 1 and 2 |
| 4. | $(p \rightarrow (q \rightarrow p))$ | Instance of axiom 1 |

Example

- | | | |
|----|---|-----------------------|
| 1. | $p \rightarrow ((q \rightarrow p) \rightarrow p)$ | Instance of axiom 1 |
| 2. | $(p \rightarrow ((q \rightarrow p) \rightarrow p)) \rightarrow ((p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p))$ | Instance of axiom 2 |
| 3. | $(p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p)$ | MP from steps 1 and 2 |
| 4. | $(p \rightarrow (q \rightarrow p))$ | Instance of axiom 1 |
| 5. | $p \rightarrow p$ | MP from steps 4 and 3 |

Example

- | | | |
|----|---|-----------------------|
| 1. | $p \rightarrow ((q \rightarrow p) \rightarrow p)$ | Instance of axiom 1 |
| 2. | $(p \rightarrow ((q \rightarrow p) \rightarrow p)) \rightarrow ((p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p))$ | Instance of axiom 2 |
| 3. | $(p \rightarrow (q \rightarrow p)) \rightarrow (p \rightarrow p)$ | MP from steps 1 and 2 |
| 4. | $(p \rightarrow (q \rightarrow p))$ | Instance of axiom 1 |
| 5. | $p \rightarrow p$ | MP from steps 4 and 3 |

Hence, $p \rightarrow p$ is a theorem.

Example



Mrs White is guilty. *w*

Miss Scarlet is guilty. *s*

Colonel Mustard is guilty. *m*

Example



Mrs White is guilty. w

Miss Scarlet is guilty. s

Colonel Mustard is guilty. m

- ▶ At least one of them is guilty.
- ▶ Not all of them are guilty.
- ▶ If Mrs White is guilty, then Colonel Mustard helped her.
- ▶ If Miss Scarlet is innocent then so is Colonel Mustard.

Example



Mrs White is guilty. w

Miss Scarlet is guilty. s

Colonel Mustard is guilty. m

- ▶ At least one of them is guilty.
- ▶ Not all of them are guilty.
- ▶ If Mrs White is guilty, then Colonel Mustard helped her.
- ▶ If Miss Scarlet is innocent then so is Colonel Mustard.

$$w \vee s \vee m$$

$$\neg(w \wedge s \wedge m)$$

$$w \rightarrow m$$

$$\neg s \rightarrow \neg m$$

The questions

Define

$$\Phi := \{w \vee s \vee m, \neg(w \wedge s \wedge m), w \rightarrow m, \neg s \rightarrow \neg m\}$$

The questions

Define

$$\Phi := \{w \vee s \vee m, \neg(w \wedge s \wedge m), w \rightarrow m, \neg s \rightarrow \neg m\}$$

- $\Phi \models s$?

The questions

Define

$$\Phi := \{w \vee s \vee m, \neg(w \wedge s \wedge m), w \rightarrow m, \neg s \rightarrow \neg m\}$$

- $\Phi \models s$?
- $\Phi \models \neg w$?

The questions

Define

$$\Phi := \{w \vee s \vee m, \neg(w \wedge s \wedge m), w \rightarrow m, \neg s \rightarrow \neg m\}$$

- $\Phi \models s$?
- $\Phi \models \neg w$?
- $\Phi \models m$?

The questions

Define

$$\Phi := \{w \vee s \vee m, \neg(w \wedge s \wedge m), w \rightarrow m, \neg s \rightarrow \neg m\}$$

- $\Phi \models s$?
- $\Phi \models \neg w$?
- $\Phi \models m$?
- $\Phi \models \neg m$?

Updates

Which are the possibilities?

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

~~$\{\}$~~ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **second** premise $\neg(w \wedge s \wedge m)$ is false:

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **second** premise $\neg(w \wedge s \wedge m)$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ ~~$\{w, s, m\}$~~

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **second** premise $\neg(w \wedge s \wedge m)$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{\cancel{w, s, m}\}$

Remove cases where the **third** premise $w \rightarrow m$ is false:

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **second** premise $\neg(w \wedge s \wedge m)$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ ~~$\{w, s, m\}$~~

Remove cases where the **third** premise $w \rightarrow m$ is false:

$\{\}$ ~~$\{w\}$~~ $\{s\}$ $\{m\}$ ~~$\{w, s\}$~~ $\{w, m\}$ $\{s, m\}$ ~~$\{w, s, m\}$~~

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **second** premise $\neg(w \wedge s \wedge m)$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{\cancel{w, s, m}\}$

Remove cases where the **third** premise $w \rightarrow m$ is false:

$\{\}$ $\{\cancel{w}\}$ $\{s\}$ $\{m\}$ $\{\cancel{w, s}\}$ $\{w, m\}$ $\{s, m\}$ $\{\cancel{w, s, m}\}$

Remove cases where the **fourth** premise $\neg s \rightarrow \neg m$ is false:

Updates

Which are the possibilities?

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **first** premise $w \vee s \vee m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **second** premise $\neg(w \wedge s \wedge m)$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **third** premise $w \rightarrow m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Remove cases where the **fourth** premise $\neg s \rightarrow \neg m$ is false:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w, s\}$ $\{w, m\}$ $\{s, m\}$ $\{w, s, m\}$

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$?

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$? **Yes!**

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$? **Yes!**
- $\Phi \models \neg w$?

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$? **Yes!**
- $\Phi \models \neg w$? **Yes!**

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$? **Yes!**
- $\Phi \models \neg w$? **Yes!**
- $\Phi \models m$?

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$? **Yes!**
- $\Phi \models \neg w$? **Yes!**
- $\Phi \models m$? **No!**

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$? **Yes!**
- $\Phi \models \neg w$? **Yes!**
- $\Phi \models m$? **No!**
- $\Phi \models \neg m$?

Updates

Only the following possibilities make **all** premises in Φ true:

$\{\}$ $\{w\}$ $\{s\}$ $\{m\}$ $\{w,s\}$ $\{w,m\}$ $\{s,m\}$ $\{w,s,m\}$

- $\Phi \models s$? **Yes!**
- $\Phi \models \neg w$? **Yes!**
- $\Phi \models m$? **No!**
- $\Phi \models \neg m$? **No!**